


# Effect of Fragmentation Models on Atmospheric Energy Deposition

## Paul J. Register, Donovan L. Mathias

**NASA Advanced Supercomputing Division** NASA Ames Research Center, Moffett Field, CA 94035, USA


July 7-9 Workshop at NASA Ames




### **OBJECTIVE & APPROACH**

The ultimate goal of our work is to understand the risk posed by asteroids with uncertain entry and strength parameters. During atmospheric entry, bolides deposit energy as they fragment, ablate, and are subjected to drag forces. Simplified fragmentation modeling approaches found in existing literature predict different energy deposition curves and resulting blast footprints. We compare four models to assess their fidelity, sensitivities to entry and strength parameters, and effects on blast wave risk estimates. We modeled energy deposition as a function of altitude, compared results against the Chelyabinsk meteoroid case and a high-fidelity simulation, and used a blast wave propagation model to compute resulting ground damage footprints. While the physics of asteroid trajectory, ablation, and fragmentation are important, extracting the key features and sensitivities of the entry process into simpler simulations is vital to producing an effective risk model.


#### **MODELS** Liquid Drop/"Pancake" **Collective Wake** (Hills & Goda, 1993) (Revelle, 2005) Ram pressure exceeds bolide Ram pressure exceeds bolide strength. strength. Debris cloud forms with twice the previous Debris cloud forms drag area and increased strength. Mass breaks again. Effective mass remains Bolide mass "pancakes" from constant while area doubles. increased drag. Mass continues to flatten Process continues to ground, under common bow shock always under common bow shock. until end of flight. **Combination Model – Progressive Fragmentation – Progressive Fragmentation with "Pancaking" Non-Collective Wake** (Mehta et al., PDC 2015) Ram pressure exceeds bolide strength. Debris cloud forms and "pancakes" Ram pressure exceeds bolide strength. Bolide under a common bow shock. breaks into two children Cloud reaches a critical radius and of nearly equal mass. separates into two fragments of nearly equal mass. Fragments obtain separate Children have bow shocks and are then treated greater strength, independently. but continue to Pressure exceeds increased fragment as ram fragment strengths again, forming pressure increases. additional debris clouds. Process continues to ground. Pieces Process continues to ground. Bolide pieces eventually cease fragmentation and eventually cease fragmentation and may slow to may slow to terminal velocity. terminal velocity.




appears to be capable of capturing key features (such as




- Strength = Ram pressure at break point  $\rightarrow \sigma = \rho_{air} v^2$ .
- $v = f(\rho_{air})$  so strength can be prescribed based on desired break altitude.
- α varies with the pre-cracked nature of the asteroid's internal structure, though the fracture pattern may change as the bolide fragments.
- Assuming constant  $\alpha$ , a higher value results in fewer and more pronounced fragmentations.
- Shows limitation of pancake model in allowing for existing structure of bolide.





fragmentation



## **CONCLUSIONS & FUTURE WORK**

Although they are at opposite extremes in terms of complexity, the pancake and combination models produce similar curves in all of the above studies and seem to describe the airburst process more intuitively than other existing models. Used in tandem, these two models can roughly outline the energy deposition and relevant features of the asteroid entry process. Future work includes refining both asteroid breakup and blast wave models, as well as simulating additional observed impact events.

## **REFERENCES**

Mehta, P. M., Minisci, E. M., Vasile, M., Break-up Modelling and Trajectory Simulation Under Uncertainty for Asteroids, 4th IAA PDC (Italy), 2015.

Hills, J. G., Goda, M. P., The Fragmentation of Small Asteroids in the Atmosphere, The Astronomical Journal (105-3), 1114-1144, 1993.

Popova, O. P., et al., Very low strengths of interplanetary meteoroids and small asteroids, Meteoritics & Planetary Science (46-10), 1525, 2011.

• Popova, O. P., Jenniskens, P., Emel'yanenko, V., et al., Chelyabinsk Airburst, Damage Assessment, Meteorite Recovery, and Characterization, Science (342), 1069, 2013. • Revelle, D. O., Recent Advances in Bolide Entry Modeling: A Bolide Potpourri, Earth, Moon, and Planets (95), 441-476, 2004.